



IMC9

THE BIOLOGY OF FUNGI

Edinburgh, UK • 1– 6 August 2010

Photo of Edinburgh courtesy of Scottish Viewpoint

Programme Book



www.imc9.info

Hosted by



British Mycological Society



Administration by



P3.139	Direct molecular identification of dothideomycete specimens on plants using PCR-TGGE analysis T. Mikawa*, S. Endo, M. Hasegawa, F. Ikeda, K. Hamano, N. Kinjo
P3.140	Species of podosporea recovered from the dung of herbivores and their ecology J. K. Misra*
P3.141	A new coelomycetous genus associated with die-back symptoms on apple trees in South Africa M. Cloete, J. de Gruyter, P.H. Fourie, P.W. Crous, L. Mostert*
P3.142	Phylogeny of the genus Bryoria (Lecanoromycetes, Ascomycota) L. Myllys*, S. Velmala, H. Holien, T. Goward, P. Halonen
P3.143	On a Zygosporium species isolated from a termite nest with view of the generic reevaluation G. Okada*, K-D An, Y. Aramsirirujwet, M. Ohkuma
P3.144	A multigene phylogenetic assessment of species boundaries in <i>Otidea</i> I. Olariaga*, K. Hansen
P3.145	Diversity and phylogeny of Lecanoraceae (Lecanorales: Ascomycota) in SE Asia K. Papong*, K. Boonpragob, H.T. Lumbsch
P3.146	Multi-locus analysis of a citreoviridin-producing isolate previously identified as <i>Penicillium</i> NRRL 13013 S.W. Peterson*, Z. Jurjevic, J. C. Frisvad
P3.147	Two new species of fungal endophytes colonise Pinus sylvestris needles in Scotland S.N.A. Reignoux*, S. Green, R.A. Ennos
P3.148	Integrated morphologic, metabolic, molecular and spectral methods for the identification of <i>Aspergillus</i> Section <i>Flavi</i> isolated from Portuguese almonds P. Rodrigues*, C. Santos, A. Venâncio, N. Lima
P3.149	Taxonomic analysis, a first step for the promotion of the safe use of fungi in cheese fermentations. J. Ropars*, J. Dupont, P. Renault
P3.150	Getting to know the fungi in Soil Clone Group 1 A. Rosling*, K. Cruz Martinez, A. Menkis, K. Ihrmark, S. Holmström, A. Broberg et al
P3.151	High levels of Botryosphaeriaceae diversity on native and introduced Acacia spp. in South Africa J.A. Van der Linde, B.A.D. Begoude, J. Roux*
P3.152	New insights into systematics of lichenicolous fungi C. Ruibal*, A.M. Millanes, D.L. Hawksworth
P3.153	Biodiversity of ascomycetes on trees native to the Andean Patagonian forests in Argentina R.M. Sánchez*, M.V. Bianchinotti
P3.154	A new species of pestalosphaeria M.E. barr on bamboo from Thailand N. Sanoamuang*
P3.155	Re-discovery of type specimens of <i>Cordyceps</i> and its allies described by Dr. Yosio Kobayasi H. Sato*, S. Ban, H. Masuya, T. Hosoya
P3.156	Species of nephroma endemic to macaronesia belong to two strongly supported clades each comprizing a widespread holarctic species and all are of recent origin E. Sérusiaux*, J. C. Villarreal, T. Wheeler, B. Goffinet
P3.157	Patterns of host specificity in genus Abrothallus (lichenicolous Ascomycota) A. Suija*, S. Pérez-Ortega, A. Crespo
P3.158	Four new species of Fusarium described from natural ecosystems in Australia M.H. Laurence, J.L. Walsh, D.M. Robinson, E.C.Y. Liew, L.W. Burgess, B.A. Summerell*
P3.159	Phylogeny and evolution of the tribe Erysipheae (Erysiphaceae) with special reference to appendage morphology and host plants S. Takamatsu*, Y. Shiroya, S. Limkaisang, H. Ito, V. Heluta
P3.160	Phylogenetic and geographic distribution of the mating type idiomorphs in aurofusarin-producing Hypomyces spp. H. Tamm*, K. Pöldmaa
P3.161	Molecular phylogeny of two coelomycetous genera with stellate conidia, <i>Prosthemium</i> and <i>Asterosporium</i> K. Tanaka*, V.A. Mel'nik, M. Kamiyama, K. Hirayama, T. Shirouzu
P3.162	Genomic and metabolomic analyses as tools for <i>Fusarium</i> classification U. Thrane*
P3.163	Evolution analysis of yeast Saccharomyces cerevisiae based on rDNA ITS region C.J. Tien*, P.H. Wang
P3.164	Phylogenetic and vegetative compatible grouping analyses of <i>Sclerotinia homoeocarpa</i> causing dollar spot of turf grasses in Japan Takao Tsukiboshi*, Toshihiro Hayakawa, Ikuko Okabe, Koya Sugawara
P3.165	Fungi of Serbia and the Western Balkans G. Milosevic, D. Krstajic, B. Uzelac*
P3.166	The lichen genus Caloplaca Th. Fr. (Teloschistaceae) in Chile R. Vargas*
P3.167	The biogeographic patterns in pyrenomycetous fungi and their taxonomy

Integrated morphologic, metabolic, molecular and spectral methods for the identification of *Aspergillus* Section *Flavi* isolated from Portuguese almonds

P. Rodrigues*^{1,2}, C. Santos¹, A. Venâncio¹, N. Lima¹

¹University of Minho, Portugal, ²CIMO-Mountain Research Centre, Portugal

Aspergillus is a large genus, with a complex and ever evolving taxonomy. Section *Flavi* is one of the most significant Sections in the Genus. Taxonomy and species identification is subject of great interest for scientists aiming to clarify the species concept and limits within the section. Furthermore, this Section comprises both toxigenic and non-toxigenic species/strains, with great interest to food industry.

Taxonomy of Section *Flavi* currently depends on multivariate approaches entailing both phenotypic (morphology, extrolite profiles) and molecular traits. No method works flawlessly in recognising species, so polyphasic approaches should be given emphasis in taxonomic decision-making.

This work aims to identify strains from Section *Flavi* by combining various methods, including morphological, biochemical and molecular ones, as well as the novel approach based on spectral analysis by MALDI-TOF-ICMS.

About 350 fungal isolates from Portuguese almonds, all belonging to Section *Flavi*, were characterised morphologically and biochemically. From those, 26 were selected for further analysis. Mycotoxigenic patterns, partial calmodulin gene sequences and MALDI-TOF protein spectra were analysed, dendrograms were created and results were compared.

We obtained good agreement between methods on species level identification. Additionally, the few incongruencies detected between morphological and biochemical data were well resolved with the molecular and spectral analyses.

Keywords: *Aspergillus* Section *Flavi*, taxonomy, polyphasic identification